

Geomorphometric analysis of river basins of the Volga Federal District using SRTM and Aster GDEM data

Ivanov M., Yermolaev O.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

A spatial database of geomorphometric parameters of 1:200 000 scale based on river basin approach has been created for the Volga Federal District for the first time. Watersheds created with semi-automated method of terrain and hydrological modeling in TAS GIS and WhiteBox GIS were used as spatial units. DEMs SRTM and Aster GDEM and hydrographic network vectorized from topographic maps were used as input data. Basic morphometric characteristics of relief such as mean height, slope, slope length, height range, river network density and factor LS were calculated for each watershed using the above-mentioned DEM. From zonal statistics for river basins, the average values of these characteristics were obtained. Spatial analysis and interpretation of morphometric characteristics of relief based on existing scales and classifications was carried out. The accuracy of the results was assessed by comparing with characteristics calculated by 100m-resolution DEM based on topographic map of scale 1:50000. Small average errors obtained prove reliability of the results.

<http://dx.doi.org/10.21046/2070-7401-2017-14-2-98-109>

Keywords

ASTER GDEM, Relief morphometry, SRTM, Volga Federal District

References

- [1] Desmet P.J.J., Govers G. A GIS procedure for automatically calculating the USLE LS factor on topographically complex landscape units//Journal of Soil and Water Conservation. 1996. Vol. 51. Issue 5. pp. 427-433.
- [2] Ermolaev O.P., Mal'tsev K.A., Ivanov M.A. Automated Construction of the Boundaries of Basin Geosystems for the Volga Federal District//Geography and Natural Resources. 2014. Vol. 35. No. 3. pp. 222-228.
- [3] Kinnell P.I.A. Alternative approaches for determining the USLE-M slope length factor for grid cells//Soil Science Society America Journal. 2005. Issue 69. pp. 674-680.
- [4] Lindsay J.B. 2014 The Whitebox Geospatial Analysis Tools project and open-access GIS//Proceedings of the GIS Research UK 22nd Annual Conference, The University of Glasgow. 2014. pp. 16-18.
- [5] Maltsev K.A., Yermolaev O.P., Mozzherin V.V. Suspended sediment yield mapping of Northern Eurasia//Proceedings IAHS. 2015. pp. 326-332.
- [6] Maltsev K., Yermolaev O., Mozzherin V. Mapping and spatial analysis of suspended sediment yields from the Russian Plain//Proceedings IAHS-AISH. 2012. pp. 251-258.
- [7] Moore I.D., Grayson R.B., Ladson A.R. Digital terrain modelling: a review of hydrological, geomorphological, and biological applications//Hydrological Processes. 1991. Vol. 5. Issue 1. pp. 3-30.

- [8] O'Callaghan J.F., Mark D.M. The extraction of drainage networks from digital elevation data//Computer Vision, Graphics and Image Processing. 1984. Issue 28. pp. 323-344.
- [9] Panagos P., Borrelli P., Meusburger K. A new European slope length and steepness factor (LS-Factor) for modeling soil erosion by water//Geosciences. 2015. Issue 5. pp. 117-126.
- [10] Wischmeier W.H., Smith D.D. Predicting rainfall erosion losses-A guide to conservation planning. Agriculture Handbook. 1978. No. 537. US Department of Agriculture, Washington DC.
- [11] Yermolaev O.P., Usmanov B.M., Muharamova S.S. The basin approach and mapping to the anthropogenic impact assessment on the east of the Russian plain//International Journal of Applied Engineering Research. 2015. Vol. 10. Issue 20. pp. 41178-41184.
- [12] Ermolaev O.P., Mal'tsev K.A., Ispol'zovanie tsifrovyykh modelei rel'efa dlya avtomatizirovannogo postroeniya granits vodosborov (Using dems for automatic plotting of catchments), Geomorfologiya, 2014, No. 1, pp. 45-52.
- [13] Ermolaev O.P., Mal'tsev K.A., Mozzherin V.V., Mozzherin V.I., Global'naya geoinformatsionnaya sistema "Stok vzveshennykh nanosov v rechnykh basseynakh Zemli" (Global geoinformation system "Suspended Sediment Yield in the River Basins of the Earth"), Geomorfologiya, 2012, No. 2, pp. 50-58.
- [14] Zaslavskii M.N., Eroziya pochv. (Soil erosion), Moscow: Mysl', 1979, 245 p.
- [15] Zaslavskii M.N., Kashtanov A.N. Pochvozashchitnoe zemledelie (Conservation agriculture.), Moscow: Rossel'khozizdat, 1979, 207 p.
- [16] Kiryushin V.I., Teoriya adaptivno-landshaftnogo zemledeliya i proektirovaniya agrolandshaftov (The theory of adaptive-landscape agriculture and agricultural landscapes design), Moscow: Kolos, 2011, 443 p.
- [17] Chentsov V.N., Morfometricheskie pokazateli na geomorfologicheskoi karte melkogo masshtaba (The morphometric indicators on small scale geomorphological map), Trudy In-ta geografii AN SSSR, 1948, Vol. 39, pp. 291-306.
- [18] Desmet P.J.J., Govers G., A GIS procedure for automatically calculating the USLE LS factor on topographically complex landscape units, Journal of Soil and Water Conservation, 1996, Vol. 51, No. 5, pp. 427-433.
- [19] Ermolaev O.P., Mal'tsev K.A., Ivanov M.A., Automated Construction of the Boundaries of Basin Geosystems for the Volga Federal District, Geography and Natural Resources, 2014, Vol. 35, No. 3, pp. 222-228.
- [20] Kinnell P.I.A., Alternative approaches for determining the USLE-M slope length factor for grid cells, Soil Science Society America Journal, 2005, No. 69, pp. 674-680.
- [21] Lindsay J.B., 2014 The Whitebox Geospatial Analysis Tools project and open-access GIS, Proceedings of the GIS Research UK 22nd Annual Conference, The University of Glasgow, 2014, pp. 16-18.
- [22] Maltsev K.A., Yermolaev O.P., Mozzherin V.V., Suspended sediment yield mapping of Northern Eurasia, Proceedings IAHS, 2015, pp. 326-332.
- [23] Maltsev K., Yermolaev O., Mozzherin V., Mapping and spatial analysis of suspended sediment yields from the Russian Plain, Proceedings IAHS-AISH, 2012, pp. 251-258.
- [24] Moore I.D., Grayson R.B., Ladson A.R., Digital terrain modelling: a review of hydrological, geomorphological, and biological applications, Hydrological Processes, 1991, Vol. 5, No. 1, pp. 3-30.
- [25] O'Callaghan J.F., Mark D.M., The extraction of drainage networks from digital elevation data, Computer Vision, Graphics and Image Processing, 1984, No. 28, pp. 323-344.
- [26] Panagos P., Borrelli P., Meusburger K., A new European slope length and steepness factor (LS-Factor) for modeling soil erosion by water, Geosciences, 2015, No. 5, pp. 117-126.
- [27] Wischmeier W.H., Smith D.D., Predicting rainfall erosion losses-A guide to conservation planning. Agriculture Handbook, US Department of Agriculture, Washington DC, 1978, No. 537.
- [28] Yermolaev O.P., Usmanov B.M., Muharamova S.S., The basin approach and mapping to the anthropogenic impact assessment on the east of the Russian plain, International Journal of Applied Engineering Research, 2015, Vol. 10, No. 20, pp. 41178-41184.